## **RAMIK**<sup>®</sup> Brown

(EPA REG. NO. 2393-497)

### SUPPLEMENTAL LABELING

### FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MONTANA

### WEATHER-RESISTANT RODENTICIDE FOR CONTROL OF ORCHARD MICE

### **ACTIVE INGREDIENTS:**

Diphacinone (2-Diphenylacetyl-1,3-

 Indandione
 0.005%

 INERT INGREDIENTS:
 99.995%

 Total
 100.000%

EPA SLN No MT-790022 EPA REG No 2393-497

# CAUTION KEEP OUT OF REACH OF CHILDREN

Keep out of reach of children, pets, wildlife, and domestic animals.

Keep out of lakes, streams or ponds.

Wear rubber gloves when handling this product. If swallowed by human beings, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of vitamin  $K_1$ , combined with blood transfusions, is indicated as in the case of hemorrhage caused by overdose of bishydrooxycoumarin (Dicumarol).

## IN ALL CASES OF HUMAN INGESTION IMMEDIATELY NOTIFY A PHYSICIAN.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of pesticide application

Bearing and Non-bearing Orchards of Tree Fruits (Stone & Pome) and Nuts.

ORCHARD	RATE/ACRE	LIMITATIONS
MICE Symfo as Famasina	OF RAMIK	/ DIRECTIONS
Surface Foraging	Aerial or	Dormant
Voles	Ground	Application
	Broadcast	Only
	Treatment	Do not graze
	Apply 20 lbs of	meat or dairy
	pellets per acre	cattle in treated
	in two (2)	fields.
	applications of	Do not cut hay

Subsurface Root Feeding Voles  In-Tunnel Bait Treatment Applications of pellets per acre in two (2) applications can be made 20 to 40 days apart.  Subsurface Root Feeding Voles  In-Tunnel Bait Treatment Apply 20 lbs of pellets per acre in two (2) applications of 10 lbs each.  Applications can be made 20 to 40 days apart.  WHERE A MIXED POPULATION IS PRESENT, APPLY AS DIRECTED FOR SUBSURFACE ROOT FEEDING VOLES.  Restore treated areas and feed or use for bedding.  Restore treated areas and feed or use for bedding.  Application is not to be made until all crops are picked up.  Dormant Application Only  For each application make at least two (2) bait placements per tree, by dispersing ½ lb of Ramik Pellets per tree into active vole trails and holes (40 tree/acre planting).  Restore treated areas with thatch or other vegetative debris.  Do not graze meat or dairy cattle in treated fields.  Do not cut hay from treated areas and feed or use for bedding.  Application is not to be made until all crops are harvested and all harvestable drops are picked				
Feeding Voles  Treatment Apply 20 lbs of pellets per acre in two (2) applications of 10 lbs each. Applications can be made 20 to 40 days apart.  WHERE A MIXED POPULATION IS PRESENT, APPLY AS DIRECTED FOR SUBSURFACE ROOT FEEDING VOLES.  WOLES.  Application make at least two (2) bait placements per tree, by dispersing ½ lb of Ramik Pellets per tree into active vole trails and holes (40 tree/acre planting).  Restore treated areas with thatch or other vegetative debris.  Do not graze meat or dairy cattle in treated fields.  Do not cut hay from treated areas and feed or use for bedding.  Application application make at least two (2) bait placements per tree, by dispersing ½ lb of Ramik Pellets per tree into active vole trails and holes (40 tree/acre planting).  Restore treated areas with thatch or other vegetative debris.  Do not graze meat or dairy cattle in treated fields.  Do not cut hay from treated areas and feed or use for bedding.  Application is not to be made until all crops are harvested and all harvestable		Applications can be made 20 to 40	areas and feed or use for bedding.  Application is not to be made until all crops are harvested and all harvestable drops are picked	
up.		Treatment Apply 20 lbs of pellets per acre in two (2) applications of 10 lbs each. Applications can be made 20 to 40 days apart. WHERE A MIXED POPULATION IS PRESENT, APPLY AS DIRECTED FOR SUBSURFACE ROOT FEEDING	Application Only  For each application make at least two (2) bait placements per tree, by dispersing ½ lb of Ramik Pellets per tree into active vole trails and holes (40 tree/acre planting).  Restore treated areas with thatch or other vegetative debris.  Do not graze meat or dairy cattle in treated fields.  Do not cut hay from treated areas and feed or use for bedding.  Application is not to be made until all crops are harvested and all harvestable drops are picked	

This specimen label is intended for use only as a guide in providing general information regarding the directions, warning and cautions associated with the use of this product. As with any product, always follow the label instructions on the package before using.

### **IMPORTANT**

For best results, keep this product in a closed container, free from odors which may contaminate the bait and reduce acceptability.

Rodents may feed several times before the effects are noticed. Several days may be required for death to occur.

### STORAGE AND DISPOSAL

**PROHIBITIONS:** Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in original container in dry, cool area separately from fertilizer, feed, or foodstuffs and away from products with strong odors. Spillage should be carefully swept up and collected for disposal.

**PESTICIDE DISPOSAL:** Pesticide wastes may be toxic. Improper disposal of excess pesticide is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse plastic pails. For paper or plastic bags, or for fiber drums with liners, completely empty into application equipment by shaking and tapping container. Offer pails or drums for recycling or reconditioning or puncture before disposal. Dispose of bags, liners, or punctured containers in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**NOTICE:** Because Manufacturer and Seller have no control over storage, handling, and conditions of use, which are of critical importance, Manufacturer and Seller make no representation or warranty, either express or implied, for results due to misuse, improper handling or storage of this material. Nor do Manufacturer and Seller assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.

24(c) Registrant: Manufactured for: HACO, Inc. PO Box 7190 Madison, WI 53707

EPA SLN No. MT-790022 16310-MT (24c) Revised (3)